

The light pseudoscalar Higgs boson in NMSSM

Abdesslam Arhrib^a, Kingman Cheung^b, Tie-Jium Hou^b, Kok-Wee Song^b

^a *Faculté des Sciences et Techniques B.P 416 Tangier, Morocco*

^b *Department of Physics and NCTS, National Tsing Hua University, Hsinchu, Taiwan*

Abstract

We study the associated production of a very light pseudoscalar Higgs boson with a pair of charginos. The novel signature involves a pair of charged leptons from chargino decays and a pair of photons from the pseudoscalar Higgs boson decay, plus large missing energy at the LHC and ILC. The signal may help us to distinguish the NMSSM from MSSM, provided that the experiment can resolve the two photons from the decay of the pseudoscalar Higgs boson.

The little hierarchy problem has motivated a number of solutions to relieve the problem. One of these is to add additional singlet fields to the minimal supersymmetric standard model (MSSM). The minimal version of the latter is realized by adding a singlet Higgs field to the MSSM, and becomes the next-to-minimal supersymmetric standard model (NMSSM). It has been shown [1] that in some corners of the parameter space, the Higgs boson can decay into a pair of very light pseudoscalars such that the LEP2 limit can be evaded. The NMSSM is in fact well motivated as it provides an elegant solution to the μ problem in SUSY. Explicitly, the superpotential of the NMSSM is given by

$$W = \mathbf{h}_u \hat{Q} \hat{H}_u \hat{U}^c - \mathbf{h}_d \hat{Q} \hat{H}_d \hat{D}^c - \mathbf{h}_e \hat{L} \hat{H}_d \hat{E}^c + \lambda \hat{S} \hat{H}_u \hat{H}_d + \frac{1}{3} \kappa \hat{S}^3. \quad (1)$$

The Higgs sector of the NMSSM consists of the usual two Higgs doublets H_u and H_d and an extra Higgs singlet S . With the additional singlet Higgs field, there are one more CP-even and one more CP-odd Higgs bosons, and one more neutralino than those in the MSSM. One particular feature of the NMSSM is the allowable light pseudoscalar boson A_1 , which is consistent with existing data. Since this A_1 mainly comes from the singlet Higgs field, it can escape all the experimental constraints when the mixing angle with the MSSM Higgs fields goes to zero. It was pointed out in Ref. [1] that even in the zero mixing limit, there is always a SM-like Higgs boson that decays into a pair of A_1 's, which helps the Higgs boson to evade the LEP bound. The possibility to detect such light pseudoscalar Higgs bosons coming from the H_1 decay was studied using the two photon mode of the A_1 , but the two photons may be too collimated for realistic detection [2]. Here we probe another novel signature in the zero-mixing limit [3]. The unique term $\lambda S H_u H_d$ in the superpotential gives rise to the coupling of $\lambda S \tilde{H}_u \tilde{H}_d$, which includes the neutral and charged Higgsinos. We study the associated production of a light pseudoscalar Higgs boson with a chargino pair in the zero-mixing limit at the LHC and ILC. Provided that the pseudoscalar is very light and the mixing angle is less than 10^{-3} , the dominant decay mode of A_1 is a pair of photons. Thus, the novel signature for the production is a pair of charged leptons and a pair of photons plus large missing energy.

The extra singlet field is allowed to couple only to the Higgs doublets of the model, the supersymmetrization of which is that the singlet field only couples to the Higgsino doublets. Consequently, the couplings of the singlet S to gauge bosons and fermions will only be manifest via their mixing with the doublet Higgs fields. After the Higgs fields take on the VEV's and rotating away the Goldstone modes, we are left with a pair charged Higgs bosons, 3 real scalar fields, and 2 pseudoscalar fields. The mass matrix for the two pseudoscalar Higgs bosons P_1 and P_2 is

$$V_{\text{pseudo}} = \frac{1}{2} (P_1 \ P_2) \mathcal{M}_P^2 \begin{pmatrix} P_1 \\ P_2 \end{pmatrix} \quad (2)$$




Figure 1: Decay branching ratios for the light pseudoscalar Higgs boson versus the mixing angle $\sin \theta_A$ for $\lambda = 1$, $\mu = 150$, $M_2 = 500$ GeV. (a) $m_{A_1} = 0.1$ GeV and (b) $m_{A_1} = 5$ GeV.

with

$$\begin{aligned}
\mathcal{M}_{P_{11}}^2 &= M_A^2, \quad \mathcal{M}_{P_{12}}^2 = \mathcal{M}_{P_{21}}^2 = \frac{1}{2} \cot \beta_s (M_A^2 \sin 2\beta - 3\lambda \kappa v_s^2), \\
\mathcal{M}_{P_{22}}^2 &= \frac{1}{4} \sin 2\beta \cot^2 \beta_s (M_A^2 \sin 2\beta + 3\lambda \kappa v_s^2) - \frac{3}{\sqrt{2}} \kappa A_\kappa v_s, \\
M_A^2 &= \frac{\lambda v_s}{\sin 2\beta} (\sqrt{2} A_\lambda + \kappa v_s),
\end{aligned} \tag{3}$$

and $\tan \beta = v_u/v_d$ and $\tan \beta_s = v_s/v$ and $v^2 = v_u^2 + v_d^2$. The pseudoscalar fields are further rotated to the diagonal basis (A_1, A_2) through a mixing angle:

$$\begin{pmatrix} A_2 \\ A_1 \end{pmatrix} = \begin{pmatrix} \cos \theta_A & \sin \theta_A \\ -\sin \theta_A & \cos \theta_A \end{pmatrix} \begin{pmatrix} P_1 \\ P_2 \end{pmatrix} \tag{4}$$

At tree-level the mixing angle is given by

$$\tan \theta_A = \frac{\mathcal{M}_{P_{12}}^2}{\mathcal{M}_{P_{11}}^2 - m_{A_1}^2} = \frac{1}{2} \cot \beta_s \frac{M_A^2 \sin 2\beta - 3\lambda \kappa v_s^2}{M_A^2 - m_{A_1}^2}. \tag{5}$$

In the approximation of large $\tan \beta$ and large M_A , which is normally valid in the usual MSSM, the tree-level CP-odd masses can be written as [4]

$$m_{A_2}^2 \approx M_A^2 (1 + \frac{1}{4} \cot^2 \beta_s \sin^2 2\beta), \quad m_{A_1}^2 \approx -\frac{3}{\sqrt{2}} \kappa v_s A_\kappa. \tag{6}$$

We are interested in the case that A_1 is very light. From Eq. (6) it can be seen that m_{A_1} can be very small if either κ or A_κ is very small. We can achieve a small mixing angle by the cancellation between the two terms in the numerator of Eq.(5), by setting

$$M_A^2 \sin 2\beta - 3\lambda \kappa v_s^2 = \sqrt{2} \lambda v_s (A_\lambda - \sqrt{2} \kappa v_s) \approx 0 \quad \Rightarrow A_\lambda \approx \sqrt{2} \kappa v_s. \tag{7}$$

In the limit of zero mixing, the A_1 only couples to a pair of charginos and neutralinos. Therefore, the dominant decay mode is $\gamma\gamma$ via a chargino loop if m_{A_1} is very light. When we turn on the small mixing angle, other modes, such as $q\bar{q}$, $\ell^+\ell^-$, and gg , appear, which will eventually dominate when the mixing angle is larger than $O(10^{-3})$. We show a typical




Figure 2: The differential cross section versus the sine of the opening angle between the two photons for $\lambda = 1$ and $\sin \theta_A = 10^{-4}$ at the LHC. Requirements of $p_{T_\gamma} > 10$ GeV and $|y_\gamma| < 2.6$ are imposed.

decay branching ratio versus the mixing angle for $m_{A_1} = 0.1, 5$ GeV in Fig. 1. As long as $\sin \theta_A < 10^{-3}$ the $\gamma\gamma$ dominates the decay of A_1 .

Note that the cross section is insensitive to $\sin \theta_A$ as long as it is less than 10^{-2} . Also, in this near-zero mixing region, the cross section scales as λ^2 . The signature is very spectacular with a pair of charged leptons and a pair of photons with a large missing energy. One can also increase the detection rates by including the hadronic decay of the charginos. Therefore, in the final state we can have (i) two charged leptons + two photons + $\cancel{E}_T$, (ii) one charged leptons + two jets + two photons + $\cancel{E}_T$, or (iii) 4 jets + two photons + $\cancel{E}_T$.

The critical issue here is whether the LHC experiment can resolve the two photons in the decay of the pseudoscalar Higgs boson. We perform a monte carlo study for the production of $\tilde{\chi}_1^+ \tilde{\chi}_1^- A_1$ followed by the decay of $A_1 \rightarrow \gamma\gamma$. We impose on the photons:

$$p_{T_\gamma} > 10 \text{ GeV}, \quad |y_\gamma| < 2.6, \quad (8)$$

which are in accord with the ECAL of the CMS detector [5]. The resolution of the “preshower” detector quoted in the report is as good as 6.9 mrad. We shall use 10 mrad as our minimum separation of the two photons. We show the distribution of the sine of the opening angle between the two photons for $M_{A_1} = 0.1, 1, 5$ GeV in Fig. 2. It is easy to understand that for A_1 as light as 0.1 GeV all the cross sections are within the opening angle $\theta_{\gamma\gamma} < 0.01$ rad. When M_{A_1} increases to 1 GeV, more than half of the cross sections are beyond 0.01 rad. For M_{A_1} as large as 5 GeV almost all cross sections are beyond $\theta_{\gamma\gamma} > 0.01$ rad. We show the resultant cross sections for $M_{A_1} = 0.1 - 5$ GeV with $p_{T_\gamma} > 10$ GeV, $|y_\gamma| < 2.6$, and $\theta_{\gamma\gamma} > 0.01$ rad in Table 1. For a $O(100) \text{ fb}^{-1}$ luminosity, Table I shows that it is only possible to detect $m_{A_1} > 1$ GeV.

There are other constraints on a light pseudoscalar from $g - 2$, rare K and B meson decays, $B - \bar{B}$ mixing, $B_s \rightarrow \mu^+ \mu^-$, and $\Upsilon \rightarrow A_1 \gamma$ [6, 7]. However, it is obvious that in these processes the light pseudoscalar interacts via the mixing with the MSSM pseudoscalar. Thus, in the limit of zero-mixing the constraints on the light A_1 can be easily evaded.

Acknowledgement

The research was supported by the NSC of Taiwan under Grant 94-2112-M-007-010-.

Table 1: Cross sections in fb for associated production of $\tilde{\chi}_1^+ \tilde{\chi}_1^- A_1$ followed by $A_1 \rightarrow \gamma\gamma$. The cuts applied to the two photons are: $p_{T_\gamma} > 10$ GeV, $|y_\gamma| < 2.6$, and $\theta_{\gamma\gamma} > 10$ mrad.

M_{A_1} (GeV)	Cross Section (fb)
0.1	0.0
0.3	0.0405
0.5	0.12
1	0.26
3	0.42
5	0.44

References

- [1] R. Dermisek and J. F. Gunion, arXiv:hep-ph/0502105.
- [2] B. A. Dobrescu, G. Landsberg and K. T. Matchev, Phys. Rev. D **63**, 075003 (2001)
- [3] A. Arhrib, K. Cheung, T. J. Hou and K. W. Song, arXiv:hep-ph/0606114.
- [4] D. J. Miller, R. Nevzorov and P. M. Zerwas, Nucl. Phys. B **681**, 3 (2004)
- [5] See <http://cmsdoc.cern.ch/cms/TDR/ECAL/ecal.html>.
- [6] G. Hiller, Phys. Rev. D **70**, 034018 (2004).
- [7] J. F. Gunion, D. Hooper and B. McElrath, Phys. Rev. D **73**, 015011 (2006)